

## Review Article

# From Incision to Recovery: Advances and Challenges in Wound Closure After Total Knee Arthroplasty

Mohamed Elfekky<sup>1a</sup>, Mohadese Rajaeirad<sup>2b</sup>, Emam Mohamed Ahmed<sup>3c</sup>, Mohamed Ahmed Hassan Abdou Elsayed<sup>3d</sup>, Eslam Mohammed Shalaby<sup>3e</sup>, Mohamad Aikrch Al Haroun<sup>3f</sup>, Sari Aidek<sup>3g</sup>, Farivar Bagheri<sup>4h</sup>, Hamidreza Esmaeili<sup>5i</sup>, Samih Tarabichi<sup>6j</sup>

---

a Dr. Elfekky, a celebrated best Orthopedic Surgeon in Dubai and one of the region's top experts in knee surgery in Dubai, brings over two decades of experience and international acclaim with four patents in ligament-sparing knee arthroplasty and 3D printing technology. As the Director of Tarabichi Joint Care, he is dedicated to advancing orthopedic surgery worldwide. From treating international patients to pioneering enhanced recovery protocols, Dr. Elfekky has redefined what's possible in orthopedic care.

[Visit Dr. Mohamed Elfekky](#)

[Connect with Dr. Mohamed Elfekky on LinkedIn](#)

[Conflicts of Interest Statement for Dr. Mohamed Elfekky](#)

b Mohadese Rajaeirad is a PhD candidate and university lecturer in the field of biomechanics, and in the last 5 years she has been active in the field of orthopedic biomechanics and patient specific implant designs.

[Visit Mohadese Rajaeirad](#)

[Connect with Mohadese Rajaeirad on LinkedIn](#)

[Conflicts of Interest Statement for Mohadese Rajaeirad](#)

c Dr. Emam Ahmed is an orthopedic surgery consultant working in Dubai Health.

[Conflicts of Interest Statement for Dr. Emam Ahmed](#)

d Dr. Mohamed Ahmed Hassan is an orthopedic specialist working in Dubai Health. He has more than 15 years of experience in orthopedic surgery.

[Connect with Dr. Mohamed Ahmed Hassan on LinkedIn](#)

[Conflicts of Interest Statement for Dr. Mohamed Ahmed Hassan](#)

e [Connect with Dr. Eslam Shalaby on LinkedIn](#)

[Conflicts of Interest Statement for Dr. Eslam Shalaby](#)

f Dr. Mohamad Aikrch Al Haroun is an Orthopedic and Trauma Consultant with over 15 years of clinical and practical experience in the orthopedic specialty, including trauma, pediatric orthopedics, foot and ankle, knee arthroscopy and arthroplasty, and shoulder arthroscopy. He completed a trauma fellowship in Singapore and holds both the Arab Board certification and a Master's degree in orthopedic surgery. In addition to his clinical work, he serves as a clinical instructor in orthopedic and trauma surgery at Muhammed Ben Rashid University.

[Visit Dr. Mohamad Al Haroun](#)

[Connect with Dr. Mohamad Al Haroun on LinkedIn](#)

[Conflicts of Interest Statement for Dr. Mohamad Al Haroun](#)

g Dr. Sari Aidek, MD, is an Orthopedic Specialist, working in Dubai Health Academic Corporation.

[Conflicts of Interest Statement for Dr. Sari Aidek](#)

h Dr. Farivar Bagheri is a skilled orthopedic surgeon with 15+ years of experience, top medical credentials from Iran, and advanced training in knee arthroscopy and joint replacement across Europe.

[Visit Dr. Farivar Bagheri](#)

[Connect with Dr. Farivar Bagheri on LinkedIn](#)

[Conflicts of Interest Statement for Dr. Farivar Bagheri](#)

i Hamidreza Esmaeili is an undergraduate student of Nutrition Sciences at Isfahan University of Medical Sciences, with a strong interest in clinical nutrition and evidence-based research. He has been actively involved in academic and executive roles, including educational coordination and content development for the research committee. In addition, he has experience in managing scientific websites and producing academic content, contributing to student-led research and academic activities.

[Connect with Hamidreza Esmaeili on LinkedIn](#)

[Conflicts of Interest Statement for Hamidreza Esmaeili](#)

j Dr. Samih Tarabichi is an orthopedic surgeon focused on joint replacement.

[Visit Dr. Samih Tarabichi](#)

[Conflicts of Interest Statement for Dr. Samih Tarabichi](#)

[Visit the Open Payments Data Page for Dr. Samih Tarabichi](#)

<sup>1</sup> Orthopedic Surgery, Tarabichi Joint Care, <sup>2</sup> Biomedical Engineering, University of Isfahan, <sup>3</sup> Dubai Health Authority, <sup>4</sup> Emirates Hospital Group, <sup>5</sup> Isfahan University of Medical Sciences, <sup>6</sup> Tarabichi Joint Care

Keywords: Total knee arthroplasty, wound closure techniques, surgical site infection, barbed sutures, negative pressure wound therapy.

<https://doi.org/10.60118/001c.146307>

---

## Journal of Orthopaedic Experience & Innovation

Vol. 7, Issue 1, 2026

---

Total Knee Arthroplasty (TKA) is one of the most common orthopedic procedures globally, with wound closure playing a critical role in postoperative recovery and complication prevention. Despite advances, the optimal closure technique remains debated. This narrative review evaluates current wound closure techniques in TKA highlighting their biomechanical and clinical outcomes. Barbed sutures offer operative efficiency and comparable biomechanical integrity, reducing closure time and needle-stick injuries. Staples provide rapid application but are linked to higher postoperative pain and infection in contaminated wounds. Tissue adhesives yield better cosmetic outcomes but are limited in high-tension zones. Hybrid techniques combining adhesives and sutures enhance closure strength and cosmesis. Emerging technologies, including AI-driven monitoring, smart dressings, and negative pressure wound therapy show potential but lack standardized protocols and long-term validation. Biomechanical considerations such as closure in knee flexion and multilayered approaches significantly influence healing outcomes. No single closure method is universally optimal; technique selection should be individualized. Emphasis on personalized approaches integrating mechanical stability, biological compatibility, and technological advances will guide future wound care strategies in TKA. Robust comparative trials and interdisciplinary collaboration are essential to standardize protocols and improve long-term outcomes.

### 1. INTRODUCTION

Total Knee Arthroplasty (TKA) is a widely performed orthopedic procedure aimed at relieving pain and restoring function in patients with severe knee joint diseases. Despite advancements in surgical techniques and postoperative care, wound-related complications remain a significant concern, impacting patient outcomes and healthcare resources.

Surgical site infection (SSI) is a serious complication following total knee arthroplasty (TKA), associated with prolonged hospitalization, higher morbidity, and in severe cases the need for revision procedures. The incidence of SSI after primary TKA has been reported to range from 1.1% to 2.6%<sup>1</sup>, while the 2018 International Consensus Meeting (ICM) on Musculoskeletal Infection estimated current infection rates in knee arthroplasty at 0.4–1.5%, with projections of 0.5–1.6% by 2030, underscoring the ongoing burden and the need for optimization of modifiable risk factors (Bagheri, Niknafs, Farhadi, et al. 2025; Schwarz, Parvizi, Gehrke, et al. 2019), highlighting the vulnerability of the knee region to infection. Notably, longer operative durations have been associated with an increased risk of SSIs (Teo et al. 2018).

Beyond infection risks, the biomechanics of wound healing play a pivotal role in postoperative recovery. Factors such as incision placement, closure technique, and joint movement can influence soft tissue tension, patellar tracking, and overall knee function. Improper wound healing may contribute to complications like arthrofibrosis and limited range of motion, underscoring the need for a compre-

hensive understanding of both surgical and rehabilitative strategies.

While various wound closure methods—such as sutures, staples, and tissue adhesives—are employed in TKA, there is a lack of consensus on the optimal approach. Emerging technologies, including smart dressings and AI-driven monitoring systems, offer promising avenues for enhancing wound care but require further investigation.

This narrative review aims to synthesize current knowledge on wound closure techniques in TKA, explore biomechanical considerations, evaluate postoperative management strategies, and identify gaps in the literature to guide future research.

### 2. METHODOLOGY

This narrative review synthesizes current evidence on wound closure techniques in TKA, with a focus on clinical outcomes, biomechanical factors, postoperative management, and emerging technologies. The objective was to assess best practices, identify challenges, and highlight existing research gaps in TKA wound management. To support this narrative synthesis, we conducted a comprehensive literature search across PubMed, Scopus, Web of Science, and the Cochrane Library up to April 2025. Search terms included combinations of “Total knee arthroplasty” AND (“wound closure” OR “suture techniques”), “TKA” AND (“surgical site infection” OR “wound healing”), “Barbed sutures” OR “staples” AND “TKA,” and “Negative pressure wound therapy” AND “orthopedics.” MeSH terms were applied where applicable, and additional records were identi-

fied through manual screening of reference lists. The search included English language publications comprising original research, systematic reviews, and meta-analyses relevant to wound closure, infection risk, and biomechanical outcomes. A narrative review was conducted using literature from databases including PubMed, Scopus, Web of Science, and the Cochrane Library up to April 2025. Studies included those evaluating closure methods (e.g., sutures, staples, adhesives, barbed sutures), infection rates, biomechanical properties, and rehabilitation outcomes.

### 3. CURRENT TECHNIQUES IN WOUND CLOSURE

Wound closure in total knee arthroplasty (TKA) is essential for optimal recovery, influencing complication rates, healing, and patient satisfaction (Rahbari et al. 2025). Current strategies include traditional sutures, barbed sutures, tissue adhesives, and hybrid systems, which aim to balance mechanical stability, biological compatibility, and operative efficiency while minimizing infection and wound-related complications. In the following, we discuss these closure methods, starting with conventional sutures. Traditional sutures remain the standard for precise wound approximation, though their time-consuming application and potential for wound complications warrant consideration (Khalefa, Smith, and Ahmad 2020; Eggers, Fang, and Lionberger 2011). The metal staples provide rapid closure (reducing operative time by 60-70%) but may increase postoperative pain and inflammation (Yuenyongviwat et al. 2016). Emerging technologies like absorbable staples with shape-memory properties and tissue adhesives with bacteriostatic barriers address some of these limitations while introducing new considerations, as given in [Table 1](#).

#### 3.1. EVOLUTION OF WOUND CLOSURE TECHNIQUES

The history of wound closure spans millennia, from ancient linen sutures to mid-20th-century stainless steel staples. Traditional methods prioritized mechanical strength, often at the expense of tissue reactivity and patient comfort. The advent of synthetic absorbable sutures in the 1970s marked a turning point, reducing long-term foreign body reactions (W. Zhang, Xie, and Zeng 2022). However, persistent challenges such as prolonged operative times, needle-stick injuries, and suboptimal cosmesis spurred innovation. Contemporary techniques, including barbed sutures and cyanoacrylate-based adhesives, now emphasize rapid deployment, reduced tissue trauma, and enhanced healing microenvironments (Nambi Gowri and King 2023).

#### 3.2. TRADITIONAL METHODS

**Sutures:** Conventional suturing techniques, including interrupted and continuous methods, remain the gold standard for TKA wound closure due to their precision and versatility. Absorbable polymers (polyglyconate, poligle-caprone) provide gradual strength reduction aligned with tissue healing, while non-absorbable options (polypropylene) suit high-tension areas (Nambi Gowri and King 2023;

Templeton et al. 2015). Subcuticular techniques like the buried vertical mattress minimize scarring by avoiding percutaneous knots (W. Zhang, Xie, and Zeng 2022). However, sutures require 12-15 minutes for 10 cm closures versus 3-5 minutes for staples, and their tracts increase infection risk (74% vs. 36% with staples in laparotomies) (Sundresh, Devagi, and Gopikrishna 2018).

**Staples:** Staples offer rapid wound approximation, reducing TKA closure time by 60-70% compared to sutures (Sundresh, Devagi, and Gopikrishna 2018). While efficient, they cause higher pain during removal and induce localized inflammation (Yuenyongviwat et al. 2016). However, metal staples induce localized inflammation due to persistent foreign body presence, increasing post-operative pain and wound discharge. Absorbable stapling systems, introduced in the 2010s, mitigate these issues by combining polyglycolic acid components with gradual hydrolysis. While initial failure loads for absorbable staples (54.5 N) match traditional polyglyconate sutures (56.9 N) (Templeton et al. 2015), their higher cost and limited adoption in contaminated wounds constrain widespread use (P. H. S. da Silva, Lopes, Stallmach, et al. 2023).

**Tissue Adhesives:** Cyanoacrylate adhesives (e.g., Dermabond, Prineo) provide non-invasive closure with bacterial resistance and superior cosmesis (89% vs. 54% excellent scars compared to sutures) (Dua, A, and Prasad 2021; Kulkarni et al. 2025). They demonstrate comparable wound dehiscence rates to traditional sutures (2.1% vs. 2.4%) while reducing infection rates to 1.51% (Kulkarni et al. 2025; Aravind, Sunil, Chandrasekar, et al., n.d.). Although cyanoacrylates are generally less ideal in high-tension or high-mobility regions, studies indicate that when appropriately applied, they perform effectively in TKA incisions (Sundaram et al. 2020). Limitations include exothermic reactions (45-52°C) and a 2-3% risk of allergic contact dermatitis with agents such as Dermabond, which may cause severe local reactions upon re-exposure (G. W. Lee, Kwak, and Lee 2021; Arkin and Reeder 2024; Kong et al. 2020; T. Liu, Tao, Zhao, et al. 2024; Park et al. 2021).

#### 3.3. EMERGING TECHNOLOGIES

**Barbed Sutures:** Barbed sutures provide knotless wound closure while evenly distributing tension along the incision (Nambi Gowri and King 2023). Clinical studies demonstrate their ability to reduce closure time by 23 minutes per procedure without increasing complication rates, while simultaneously decreasing needlestick injury risks (E. Li, Niu, Lu, et al. 2020). Biomechanical testing shows comparable strength to traditional sutures (54.5N vs 56.9N failure load), with both failing through tissue rupture rather than suture breakage (Templeton et al. 2015). However, they require precise implantation depth to avoid subcutaneous puckering, particularly challenging in thin-skinned patients (Nambi Gowri and King 2023).

**Absorbable Staples:** Next-generation absorbable staples made from polydioxanone (PDO) incorporate shape-memory properties that dynamically adjust tension as post-operative edema resolves. Concurrent developments include 4D-printed meshes with temperature-responsive

**Table 1. Properties and performance of TKA wound closure modalities.**

Technique	Advantages	Disadvantages	Clinical Outcomes	Biomechanical Properties
Traditional Sutures	Precise wound approximation; Versatility in technique and materials; Absorbable options align with tissue healing (Nambi Gowri and King 2023; Templeton et al. 2015)	Time-consuming (12–15 min for 10 cm incision); Infection risk through suture tracts (74%) (Sundresh, Devagi, and Gopikrishna 2018); Patient discomfort	Gold standard in many surgeries; Subcuticular sutures improve cosmetic outcomes (W. Zhang, Xie, and Zeng 2022)	Continuous sutures distribute tension evenly; Risk of peak stress at needle entry sites (Nambi Gowri and King 2023; Swain and Gupta 2015)
Metal Staples	Fast application (reduces operative time by 60–70%) (Sundresh, Devagi, and Gopikrishna 2018); Effective in high-volume procedures	Increased postoperative pain (Yuenyongviwat et al. 2016); Local inflammation; Higher infection in contaminated wounds (up to 4.7%) (Sundresh, Devagi, and Gopikrishna 2018; Lauerman, Kolesnik, Park, et al. 2018)	Shortened closure time; Less cosmetic satisfaction; Not ideal in contaminated or high-mobility wounds (Sundresh, Devagi, and Gopikrishna 2018; Lauerman, Kolesnik, Park, et al. 2018)	Stress concentrated at fixation points; May lead to edge necrosis under load (Swain and Gupta 2015)
Absorbable Staples	No need for removal; Improved comfort; Dynamic tension control with shape-memory effect	High cost; Limited use in contaminated wounds (Lu et al. 2021)	Reduced dehiscence in trials (4.1% → 1.9%)	Polydioxanone staples adjust to edema changes; Comparable initial failure load to sutures (54.5 N vs 56.9 N) (Templeton et al. 2015)
Tissue Adhesives	Rapid and non-invasive; Bacteriostatic barrier (Kulkarni et al. 2025); High cosmetic satisfaction (Dua, A, and Prasad 2021)	Risk of thermal reaction (45–52°C) (Medeiros et al. 2024); Contraindicated in high-tension/movement areas (Medeiros et al. 2024; H. G. Lee 2024)	Infection rates lower (1.51%) than sutures (2.13%) and staples (1.89%) (Kulkarni et al. 2025); 89% reported 'excellent' scar appearance (Dua, A, and Prasad 2021)	Rigid seal during early healing may hinder fibroblast migration (Medeiros et al. 2024); Not suitable under high dynamic stress (H. G. Lee 2024)
Barbed Sutures	Knotless design reduces time and needlestick risk (E. Li, Niu, Lu, et al. 2020); Even tension distribution (Nambi Gowri and King 2023)	Depth control required to prevent puckering (Nambi Gowri and King 2023)	No increase in complication rates; Operative time saved (e.g., 23 min per mastectomy); Comparable strength to traditional sutures (54.5 N) (Templeton et al. 2015)	Load distributed along wound line; Prevents peak tension points (Nambi Gowri and King 2023; Swain and Gupta 2015)
Hybrid Approaches	Combine strength and cosmetic benefit; Customized to wound type; Advanced applications (e.g., ultrasonic welding + adhesives) (H. G. Lee 2024)	Lack of standard protocols; Higher technical demands	Superior strength in models (92% of native tissue strength with adhesive-welded closure vs 85% with sutures) (H. G. Lee 2024); Promising for tracheal and joint repairs	Synergistic tension resistance and micromotion dampening (H. G. Lee 2024); Adhesives reduce epithelial shear while sutures resist gross displacement

pores that adapt to exudate volume while maintaining negative pressure therapy.

**Hybrid Methods:** Combining techniques, such as using barbed sutures for deep tissue layers and adhesives for skin closure, aims to leverage the benefits of each method (Romanini, Zanolli, Ascione, et al. 2024; Roumeliotis and Graham 2019; L. Li et al. 2023). Such approaches may optimize wound healing and cosmetic outcomes, though standard-

ized protocols remain to be established. Combining adhesives with ultrasonic suture welding exemplifies integrative approaches, with experimental models demonstrating superior strength compared to conventional methods (Romanini, Zanolli, Ascione, et al. 2024; Roumeliotis and Graham 2019; L. Li et al. 2023).

### 3.4. CLINICAL OUTCOMES AND PATIENT-CENTERED METRICS

Contemporary evidence demonstrates distinct clinical profiles for various TKA closure methods. Barbed sutures reduce operative time significantly compared to traditional techniques, with multiple studies confirming their time efficiency (Romanini, Zanolli, Ascione, et al. 2024; Khlopas et al. 2019; Eickmann and Quane 2010; Luo, Zhang, Yan, et al. 2020). Infection rates show modest variations between modalities, with meta-analyses reporting 2.13% for sutures, 1.89% for staples, and 1.51% for adhesives (Kulkarni et al. 2025). Notably, staples demonstrate higher infection rates (4.7%) in contaminated wounds, limiting their utility in non-clean cases (Sundresh, Devagi, and Gopikrishna 2018; Lauerman, Kolesnik, Park, et al. 2018). Patient-reported outcomes favor adhesives, which yield lower pain scores (1.8/10 VAS) and superior scar ratings (15-20% better than sutures) due to their non-penetrating application (Dua, A, and Prasad 2021; H. G. Lee 2024). However, adhesives show limitations in high-tension areas, where minor dehiscence (0.5-1mm) may cause local irritation (Medeiros et al. 2024). Subcuticular sutures remain a reliable option for cosmetic outcomes, though statistical significance versus alternatives varies across studies (Nepal et al. 2020; Newman et al. 2011).

### 3.5. BIOMECHANICAL PROPERTIES OF CLOSURE MATERIALS

The mechanical behavior of closure materials critically influences healing outcomes. Barbed sutures excel in tension distribution, reducing peak stress by 40-60% compared to staples through even load sharing along the wound (Nambi Gowri and King 2023; Swain and Gupta 2015). Staples concentrate stress at fixation points, potentially compromising wound margins during cyclic loading (Swain and Gupta 2015). Hybrid approaches combining sutures with adhesives offer synergistic benefits - sutures provide initial strength while adhesives minimize disruptive micromotion (H. G. Lee 2024). Material properties further impact outcomes: multifilament sutures harbor 12-15% more *S. aureus* than monofilaments due to bacterial entrapment (Sundresh, Devagi, and Gopikrishna 2018), while polypropylene non-absorbables trigger granulomatous reactions in 8-12% of patients (W. Zhang, Xie, and Zeng 2022). Adhesives avoid these issues but may restrict fibroblast migration during early healing phase (days 3-7) if overapplied (Medeiros et al. 2024). These biomechanical factors necessitate careful material selection based on surgical site demands and anticipated patient activity levels.

## 4. BIOMECHANICAL CONSIDERATIONS IN WOUND HEALING AFTER TKA

The biomechanical success of wound healing in TKA depends critically on surgical technique, soft tissue management, and postoperative rehabilitation. Closing the wound with the knee in 90° flexion, rather than full extension,

has been shown to improve early range of motion (ROM) and reduce anterior knee pain by allowing more natural soft tissue alignment under physiologic tension (Lu et al. 2021; Wang et al. 2014). Minimally invasive incisions (8-12 cm) reduce lateral skin numbness by 38% compared to traditional longer incisions (14-18 cm), preserving sensory branches of the infrapatellar nerve that contribute to proprioception (I. Alcelik, Sukeik, Pollock, et al. 2012). However, smaller incisions may increase soft tissue tension during deep flexion (>90°), potentially compromising wound integrity. In contrast, conventional midline incisions offer better surgical exposure but disrupt more vasculature, increasing dependence on collateral circulation. Optimal closure technique—such as suturing in semi-flexion (30-45°)—balances tension across the wound, reducing dehiscence risk while promoting functional recovery (Chen, Bains, Sodhi, et al. 2022). The biomechanical and clinical trade-offs of common closure methods are compared in [Table 2](#), highlighting barbed sutures' 6% complication rate versus staples' stress concentration risks.

Soft tissue tension plays a pivotal role in postoperative joint mechanics. Studies measuring intraoperative distraction forces report averages of 126N in extension and 121N in flexion, underscoring the need for precise tensioning to ensure stability and ROM (A, H, and Tj 2004; Becker, Voss, Lettner, et al. 2025). Patellar tracking is particularly sensitive to surgical technique; improper closure can alter lateral retinacular tension, increasing patellofemoral contact pressures by 12-15% during deep flexion (I. A. Alcelik et al. 2016). Meticulous attention to extensor mechanism alignment is essential to prevent maltracking, which can impair gait and accelerate wear (Gasparini, Familiari, and Ranuccio 2013). Postoperative scar formation further influences outcomes, with excessive anterior fibrosis elevating patellofemoral forces by 18% during stair ascent (Harato et al. 2016). Early mobilization—achieving >90° flexion by postoperative day 3—reduces arthrofibrosis risk by 23%, though overly aggressive rehabilitation may strain healing tissues (Chen, Bains, Sodhi, et al. 2022). Direct ice application, while reducing pain, decreases peri-incisional perfusion by 40%, potentially delaying epithelialization (Foster, Williams, Forte, et al., n.d.).

Complications arising from suboptimal wound biomechanics include patellar maltracking, arthrofibrosis, and restricted ROM ([Table 3](#)). Maltracking often results from misaligned closures or component malposition, leading to pain and functional deficits (Gasparini, Familiari, and Ranuccio 2013; Zhu et al. 2017). Arthrofibrosis, characterized by excessive scar tissue, typically stems from uncontrolled inflammation or delayed rehabilitation (Thompson, Novikov, Cizmic, et al. 2019). Physical therapy remains first-line treatment, with manipulation under anesthesia or surgical release considered for refractory cases (Wang et al. 2014; Ali, Ferguson, Singh, et al. 2023; Cheuy et al. 2017; Ipach et al. 2011). Limited ROM, influenced by preoperative stiffness and surgical trauma, can hinder basic functions like stair climbing; achieving 90° flexion early is critical for recovery (Wang et al. 2014; Ipach et al. 2011).

**Table 2. Biomechanical properties and clinical outcomes of common TKA wound closure methods**

Method	Biomechanical Impact	Clinical Outcome
<b>Barbed sutures</b>	Distribute tension evenly, reducing peak stress	6% wound complications vs. 13% with non-barbed sutures (Chen, Bains, Sodhi, et al. 2022)
<b>Staples</b>	Concentrate stress at fixation points	Higher edge necrosis risk in limited ROM patients (Dubin et al. 2022)
<b>Cyanoacrylate adhesives</b>	Rigid seals restrict micromotion early healing	2-octyl adhesive reduces wound discharge by 50% vs. non-adhesive closures (T. Liu, Tao, Zhao, et al. 2024)

**Table 3. Biomechanical triggers and clinical consequences of major TKA wound healing complications**

Complication	Biomechanical Trigger	Clinical Impact
Patellar maltracking	Altered quadriceps vector from proximal scar contracture	22% higher anterior knee pain scores (Harato et al. 2016)
Arthrofibrosis	Reduced TF joint translation from posterior capsule fibrosis	15° average flexion loss (Chen, Bains, Sodhi, et al. 2022)
Wound dehiscence	Cyclic loading exceeding suture/adhesive failure limits	2.1–4.7% revision risk (T. Liu, Tao, Zhao, et al. 2024; Chen, Bains, Sodhi, et al. 2022)

To optimize outcomes, surgeons should consider lateral-parapatellar approaches, which minimize quadriceps disruption but require careful closure to avoid sensory loss (Yuan et al. 2019; Peng et al. 2015; Shah and Shah 2024). In addition, the subvastus approach represents a minimally invasive technique that preserves the extensor mechanism, reduces postoperative pain, and facilitates faster functional recovery. Postoperative positioning also matters: supine sleeping improves terminal extension by 3° compared to lateral positions, reducing flexion contracture risk (Chen, Bains, Sodhi, et al. 2022; Cao et al. 2021). Rehabilitation timing should balance tissue healing and mobility; delaying flexion beyond 90° until days 5–7 allows fibroblast proliferation while preventing stiffness (Chen, Bains, Sodhi, et al. 2022; Foster, Williams, Forte, et al., n.d.).

## 5. POSTOPERATIVE MANAGEMENT AND WOUND HEALING CONSIDERATIONS

Early postoperative mobilization supports circulation, reduces edema, and promotes functional recovery after TKA, provided that excessive stress on the incision is avoided. Gentle range-of-motion exercises initiated within 24–48 hours and limited knee flexion ( $\leq 90^\circ$ ) during the first week help maintain tissue oxygenation and wound stability (Ghritlahre et al. 2024; Rr et al. 2019; Freccero, Van Steyn, Joslin, et al. 2022; Jaiswal et al. 2012; Mau-Moeller et al. 2014). Rehabilitation should progress gradually with careful monitoring of healing status. Adjunct measures such as manual scar mobilization or biofeedback may assist recovery when applied cautiously (Jurecka et al. 2021; Alsiri et al. 2025). Compression therapy offers mixed results and should be individualized (P. Liu et al. 2020). While postoperative rehabilitation can substantially influence wound healing outcomes, it lies beyond the primary focus of this review and is therefore only briefly addressed here.

Targeted nutritional support plays a crucial role in post-operative tissue repair. Collagen peptides provide essential amino acids that enhance collagen formation and improve scar quality (Evans and Evans 2018), while vitamin C serves as a critical cofactor for collagen synthesis and antioxidant protection (DePhillipo et al. 2018). Vitamin D supplementation, particularly high-dose preoperative regimens (300,000 IU), may reduce postoperative infections in deficient patients (Birinci, Hakyemez, Geçkalan, et al. 2024). Essential fatty acids demonstrate complex effects - omega-3s modulate inflammatory responses while omega-6s like linoleic acid promote angiogenesis (Chow and Barbul 2014; J. R. Silva et al. 2018). Additional micronutrients including zinc, vitamin A, and arginine further support immune function and cellular proliferation during wound healing (Kurdi et al. 2023; Yusuf 2022). These interventions should be individualized based on patient nutritional status and healing progression.

## 6. ADVANCED WOUND DRESSINGS AND EMERGING TECHNOLOGIES

### 6.1. MODERN WOUND DRESSINGS

Contemporary wound dressings for TKA have transitioned from basic gauze to sophisticated biomaterial systems designed to optimize healing (Table 4). Hydrocolloid dressings maintain a moist, bacteria-resistant environment that promotes natural debridement while allowing joint movement (Nguyen et al., n.d.). For wounds with moderate drainage, hydrofiber and alginate dressings provide superior fluid absorption while conforming to the surgical site (Nguyen et al., n.d.; Su et al. 2022). Clinical studies demonstrate silicone foam dressings (like Mepilex Border®) offer the best outcomes - reducing complications while improving comfort and mobility scores compared to other options (Dobbelaere et al. 2015).

**Table 4. Characteristics and Clinical Applications of Modern Wound Dressings in TKA**

Dressing Type	Key Features	Clinical Benefits in TKA	Limitations / Challenges
Hydrocolloid	Moist environment, occlusive, promotes autolytic debridement	Supports early mobilization, reduces infection risk	Limited TKA-specific data (Nguyen et al., n.d.)
Hydrofiber / Alginate	Highly absorbent, conforms to wound contours	Suitable for moderate to high exudate wounds, maintains moisture balance	May require frequent dressing changes (Nguyen et al., n.d.; Su et al. 2022)
Foam / Silicone (e.g., Mepilex®)	Cushions wound, absorbs exudate, improves comfort	Lower wound complication rates, less pain, enhanced mobility, fewer dressing changes	Higher cost in some cases (Nguyen et al., n.d.; Dobbelaere et al. 2015)
Transparent Film / Hydropolymer	Prolonged wear (up to 14 days), allows visual monitoring	High patient satisfaction, fewer dressing changes, compatibility with enhanced recovery after surgery	Limited absorption capacity (Rousseau, Plomion, and Sandy-Hodgetts 2022)
Hydrogel with bioactives	Enhances angiogenesis, reduces fibrosis, modulates healing	Faster wound healing, improved scar quality	Limited application in TKA; primarily preclinical studies (Nifontova, Safaryan, Khristidis, et al. 2024; Song, Li, Zhao, et al. 2022)
Bioengineered / Composite Dressings	Mimics ECM, includes growth factors or antimicrobials	Supports tissue regeneration, biocompatible, biodegradable	Limited clinical evidence in TKA patients (Nguyen et al., n.d.; Zarepour et al. 2024)
Smart Dressings (sensor-integrated)	Monitors temperature, pH, moisture, bacterial presence	Enables early detection, reduces hospital visits, allows real-time feedback	Experimental stage, high cost (Nguyen et al., n.d.; Tarazi et al. 2020; Jiang, Trotsyuk, Niu, et al. 2023)

Transparent hydropolymer dressings enable continuous wound monitoring with extended 14-day wear, minimizing dressing changes while maintaining visibility (Rousseau, Plomion, and Sandy-Hodgetts 2022). Cutting-edge hydrogel dressings now incorporate bioactive components such as growth factors that actively enhance tissue regeneration and scar quality (Nifontova, Safaryan, Khristidis, et al. 2024; X. Zhang, Huang, Luo, et al. 2024). The most advanced options include bioengineered scaffolds and bacterial nanocellulose dressings that mimic natural tissue while delivering antimicrobial agents, though more TKA-specific research is needed (Zarepour et al. 2024).

## 6.2. SMART MONITORING TECHNOLOGIES

Current smart wound monitoring systems (Table 5) integrate wireless sensors to track critical healing parameters including temperature, pH, moisture, and bacterial load in real-time (Jiang, Trotsyuk, Niu, et al. 2023). These technologies offer three key capabilities: (1) early complication detection through continuous monitoring, (2) remote patient tracking to reduce unnecessary clinic visits, and (3) data-driven personalization of treatment plans (Tarazi et al. 2020). While promising, their clinical adoption faces significant validation and implementation hurdles.

## 6.3. NEGATIVE PRESSURE WOUND THERAPY (NPWT)

NPWT applies controlled sub-atmospheric pressure through a sealed dressing, a technique shown to significantly improve TKA outcomes (Siqueira et al. 2016). As summarized in Table 6, the therapy enhances healing

through three key mechanisms: reduced edema, increased perfusion, and mechanical stabilization of wound edges (Song, Li, Zhao, et al. 2022; Chen, Bains, Sax, et al. 2022). Clinical studies demonstrate 40-50% lower surgical site infection rates compared to conventional dressings, particularly in high-risk patients (Song, Li, Zhao, et al. 2022).

Table 6 highlights that while NPWT's economic value grows clearer through reduced reoperations and dressing changes (Su et al. 2022; Dobbelaere et al. 2015), clinicians must balance these benefits against equipment costs and monitor for potential skin complications like blistering (Helito et al. 2017). For optimal results, NPWT should be reserved for high-risk patients (e.g., those with obesity or diabetes), while standard dressings remain appropriate for routine cases.

## 7. DISCUSSION & EXPERT COMMENTARY

### 7.1. SCIENTIFIC SYNTHESIS AND CLINICAL IMPLICATIONS

Wound closure critically influences postoperative outcomes in TKA (L. Zhang et al. 2025). Evidence favors barbed sutures for their lower complication rates (3% vs 6%) and faster closure ( $\approx 7$  minutes shorter) than traditional sutures (Chen et al. 2024; Naylor et al. 2023). Multilayer watertight closures that integrate barbed sutures with topical adhesives or polyester mesh further reduce infection and dehiscence (Snyder et al. 2021).

2-octyl cyanoacrylate adhesives enhance superficial closure outcomes, improving cosmesis, discharge control, and

**Table 5. Artificial Intelligence (AI) and Sensor-Based Wound Monitoring Technologies in TKA**

Technology	Function	Benefits for TKA Patients	Limitations / Challenges
Smart Sensor-Integrated Dressings	Real-time sensing of temperature, pH, moisture, bacterial presence	Early detection of infection or delayed healing, remote monitoring, reduced clinic visits	Still in experimental phase (Nguyen et al., n.d.; Tarazi et al. 2020; Jiang, Trotsyuk, Niu, et al. 2023)
AI Algorithms	Data analytics and complication prediction	Personalized wound care, trend analysis, alerts for early intervention	Requires validation through clinical trials (Nguyen et al., n.d.; Tarazi et al. 2020)
Closed-Loop Smart Systems	Autonomous drug delivery or electrical stimulation based on wound signals	Accelerated healing, tailored interventions	Currently under development (Jiang, Trotsyuk, Niu, et al. 2023)

**Table 6. Clinical applications of negative pressure wound therapy (NPWT) in TKA**

Feature	Clinical Impact in TKA	Benefits	Limitations / Challenges
Mechanism of Action	Applies sub-atmospheric pressure; removes exudate, promotes granulation and perfusion	Improves healing, reduces infection, enhances wound edge approximation	Requires specialized equipment and close monitoring (Song, Li, Zhao, et al. 2022; Chen, Bains, Sax, et al. 2022)
High-Risk Patient Use	Effective in patients with obesity, diabetes, or prior wound complications	Lower rates of wound dehiscence and surgical site infections	Risk of skin blistering (mitigated in newer devices) (Song, Li, Zhao, et al. 2022; Helito et al. 2017)
Economic Considerations	May reduce overall costs by preventing complications despite higher upfront cost	Fewer reoperations and dressing changes	Needs cost-benefit justification for routine implementation (Su et al. 2022; Dobbelaere et al. 2015; Helito et al. 2017)

patient satisfaction without compromising safety (T. Liu, Tao, Zhao, et al. 2024). Technique selection should remain individualized, guided by comorbidities, skin quality, and activity level (Chen et al. 2024).

Our team recommends barbed sutures for capsular closure, performed in 60–90° knee flexion to optimize alignment and recovery (Maniar, Mody, Wakankar, et al. 2023; Sanz-Ruiz et al. 2025). Triclosan-coated sutures are advised for high-risk patients, and NPWT remains beneficial in high-risk or complex cases, such as patients with obesity, diabetes, or wound-healing challenges, despite its higher cost (Ainslie-Garcia, Anderson, Bloch, et al. 2025; Krebs, Elmallah, Khlopa, et al. 2018). Emerging tools such as AI-assisted monitoring are promising but require validation. Future studies should establish standardized hybrid closure protocols and long-term cost-effectiveness analyses. [Figure 1](#) summarizes the principal domains of wound closure after TKA, including preoperative, surgical, biomechanical, and postoperative considerations.

## 7.2. EXPERT PERSPECTIVE

Based on our team's clinical experience, optimal TKA closure should balance mechanical integrity, biological compatibility, technical feasibility, and patient comfort. We recommend broader adoption of barbed sutures and advanced adhesives, which in our practice provide faster, more reliable closure than traditional techniques.


We particularly endorse flexion closure (60–90°) for improved early range of motion and reduced anterior knee pain, while recognizing its ergonomic challenges. For high-

risk patients, triclosan-coated sutures are advisable, and NPWT remains valuable in complex cases.

Emerging technologies such as absorbable staples, smart dressings, and AI monitoring hold promise but require further validation and cost-benefit assessment before routine use. We advocate for comparative effectiveness research and standardized biomechanical testing to integrate innovation into clinical protocols.

## 8. PERSISTING GAPS IN TKA WOUND CLOSURE RESEARCH

TKA closure research is limited by heterogeneous study designs and inconsistent outcome definitions, lacking high-quality comparative trials. Biomechanical insights, such as finite element modeling, rarely translate to clinical practice. Emerging technologies (absorbable staples, smart dressings, AI monitoring) face adoption challenges due to cost, limited long-term data, and insufficient randomized validation. Research is often siloed, missing interdisciplinary collaboration across surgery, rehabilitation, engineering, and nutrition. Short-term focus overlooks longitudinal outcomes like joint mobility, aesthetics, patient satisfaction, and revision risks. Prospective, multidisciplinary studies with extended follow-up are needed to address these gaps.



**Figure 1. Visual summary of key domains in wound closure after TKA, illustrating preoperative considerations, surgical closure techniques, biomechanical principles, postoperative management, and emerging technologies. Image credits: Authors. Figure created by the authors based on information synthesized from the reviewed literature.**

## 9. CONCLUSION

Wound closure in TKA is not merely a procedural step—it is central to recovery, influencing infection rates, joint function, and patient satisfaction. No universal method excels across all metrics, and technique selection should be personalized based on patient profile, tissue characteristics, and rehabilitation goals. Technologies such as barbed sutures, tissue adhesives, and hybrid methods offer benefits in operative efficiency and healing. Additionally, advanced dressings, AI-enabled monitoring, and negative pressure

therapy reflect a shift toward personalized, tech-integrated care, though they require further validation.

The future of TKA wound management lies in a multifaceted, evidence-based approach—one that unites surgical precision, mechanical insight, and patient-centered care to optimize both short- and long-term outcomes.

Submitted: August 02, 2025 EDT. Accepted: October 25, 2025 EDT. Published: May 19, 2026 EDT.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-ND-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc-nd/4.0> and legal code at <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> for more information.

## REFERENCES

- A, H., A. H., and W. Tj. 2004. "Soft-Tissue Tension Total Knee Arthroplasty." *J Arthroplasty* 19 (5). <https://doi.org/10.1016/j.arth.2004.01.003>.
- Ainslie-Garcia, M. H., L. A. Anderson, B. V. Bloch, et al. 2025. "Identifying Critical Evidence Gaps in Wound Closure and Incision Management After Total Knee Arthroplasty: Delphi Panel Insights." *J Arthroplasty* 40 (1): 111-118.e1. <https://doi.org/10.1016/j.arth.2024.06.057>.
- Alcelik, I. A., M. I. Blomfield, G. Diana, A. J. Gibbon, N. Carrington, and S. Burr. 2016. "A Comparison of Short-Term Outcomes of Minimally Invasive Computer-Assisted vs Minimally Invasive Conventional Instrumentation for Primary Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *J Arthroplasty* 31 (2): 410-18. <https://doi.org/10.1016/j.arth.2015.09.013>.
- Alcelik, I., M. Sukeik, R. Pollock, et al. 2012. "Comparison of the Minimally Invasive and Standard Medial Parapatellar Approaches for Primary Total Knee Arthroplasty." *Knee Surg Sports Traumatol Arthrosc Off J ESSKA* 20 (12): 2502-12. <https://doi.org/10.1007/s00167-012-1944-3>.
- Ali, M., C. Ferguson, I. Singh, et al. 2023. "Arthrofibrosis in Robotic Total Knee Arthroplasty: An Investigation Into How Robotic Assistance May Contribute to a Tight Knee." *JAAOS Glob Res Rev* 7 (5): e23.00025. <https://doi.org/10.5435/JAAOSGlobal-D-23-00025>.
- Alsiri, N., S. A. Alshatti, M. Al-Saffar, R. S. Bhatia, F. Fairouz, and S. Palmer. 2025. "EMMATKA Trial: The Effects of Mobilization with Movement Following Total Knee Arthroplasty in Women: A Single-Blind Randomized Controlled Trial." *J Orthop Surg* 20:181. <https://doi.org/10.1186/s13018-025-05568-8>.
- Aravind, K., C. G. Sunil, S. Chandrasekar, et al. n.d. "Cyanoacrylate vs. Sutures in Clean and Clean-Contaminated Surgical Wounds – a Randomised Control Study." *Innov Surg Sci* 9 (1): 47-54. <https://doi.org/10.1515/iss-2023-0060>.
- Arkin, L. M., and M. Reeder. 2024. "Commentary on JPOSNA® Paper, 'Dermatitis after Spine Fusion Caused by Liquid Adhesive (2-Octyl Cyanoacrylate).'" *J Pediatr Orthop Soc N Am* 7:100044. <https://doi.org/10.1016/j.jposna.2024.100044>.
- Bagheri, A., A. S. Niknafs, B. Farhadi, et al. 2025. "Incidence and Risk Factors of Surgical Site Infection After Knee Arthroplasty; a Systematic Review and Meta-Analysis." *Arch Acad Emerg Med* 13 (1): e28-e28. <https://doi.org/10.22037/aaemj.v13i1.2543>.
- Becker, R., M. Voss, J. Lettner, et al. 2025. "Mean Distraction Force Applied in Tension-Controlled Ligament-Balanced Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *Knee Surg Sports Traumatol Arthrosc* 33 (7): 2498-2526. <https://doi.org/10.1002/ksa.12629>.
- Birinci, M., Ö. S. Hakyemez, M. A. Geçkalan, et al. 2024. "Effect of Vitamin D Deficiency on Periprosthetic Joint Infection and Complications After Primary Total Joint Arthroplasty." *J Arthroplasty* 39 (9S2): S151-57. <https://doi.org/10.1016/j.arth.2024.05.012>.
- Cao, L., H. Yang, K. Sun, H. Wang, H. Fan, and W. Cheng. 2021. "The Role of Knee Position in Blood Loss and Enhancement of Recovery after Total Knee Arthroplasty." *J Knee Surg* 34 (12): 1304-9. <https://doi.org/10.1055/s-0040-1708042>.
- Chen, Z., S. S. Bains, O. C. Sax, N. Sodhi, and M. A. Mont. 2022. "Dressing Management during Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *J Knee Surg* 35:1524-32. <https://doi.org/10.1055/s-0042-1758674>.
- — —. 2024. "Optimal Method of Skin Wound Management for Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *J Knee Surg* 37 (3): 238-47. <https://doi.org/10.1055/s-0043-1768248>.
- Chen, Z., S. S. Bains, N. Sodhi, and M. A. Mont. 2022. "Outcomes of Deep Wound Management Methods During Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *Surg Technol Int* 41:289-95. <https://doi.org/10.52198/22.STI.41.OS1609>.
- Cheuy, V. A., J. R. H. Foran, R. J. Paxton, M. J. Bade, J. A. Zeni, and J. E. Stevens-Lapsley. 2017. "Arthrofibrosis Associated With Total Knee Arthroplasty." *J Arthroplasty* 32 (8): 2604-11. <https://doi.org/10.1016/j.arth.2017.02.005>.
- Chow, O., and A. Barbul. 2014. "Immunonutrition: Role in Wound Healing and Tissue Regeneration." *Adv Wound Care* 3 (1): 46-53. <https://doi.org/10.1089/wound.2012.0415>.

- DePhillipo, N. N., Z. S. Aman, M. I. Kennedy, J. P. Begley, G. Moatshe, and R. F. LaPrade. 2018. "Efficacy of Vitamin C Supplementation on Collagen Synthesis and Oxidative Stress After Musculoskeletal Injuries: A Systematic Review." *Orthop J Sports Med* 6 (10): 2325967118804544. <https://doi.org/10.1177/2325967118804544>.
- Dobbelaere, A., N. Schuermans, S. Smet, C. Van Der Straeten, and J. Victor. 2015. "Comparative Study of Innovative Postoperative Wound Dressings after Total Knee Arthroplasty." *Acta Orthop Belg* 81 (3): 454–61.
- Dua, A., N. S. A., and K. K. Prasad. 2021. "Comparison of Tissue Adhesive (N-Butyl-2-Cyanoacrylate) versus Conventional Suturing in Umbilical Hernia Surgeries." *Int Surg J* 8 (4): 1165–67. <https://doi.org/10.18203/2349-2902.isj20211291>.
- Dubin, J. A., D. Hameed, S. S. Bains, T. Board, R. Nunley, and M. A. Mont. 2022. "Effect of Various Ancillary Operating Room Techniques on Wound Healing Outcomes After Total Knee Arthroplasty." *Surg Technol Int* 43:279–88. <https://doi.org/10.52198/23.STI.43.OS1748>.
- Eggers, M. D., L. Fang, and D. R. Lionberger. 2011. "A Comparison of Wound Closure Techniques for Total Knee Arthroplasty." *J Arthroplasty* 26 (8): 1251-1258.e4. <https://doi.org/10.1016/j.arth.2011.02.029>.
- Eickmann, T., and E. Quane. 2010. "Total Knee Arthroplasty Closure with Barbed Sutures." *J Knee Surg* 23 (05): 163–68. <https://doi.org/10.1055/s-0030-1268692>.
- Evans, D. C., and B. G. Evans. 2018. "The Effects of Platelet-Rich Plasma and Activated Collagen on Wound Healing in Primary Total Joint Arthroplasty." *Orthopedics* 41 (2): e262–67. <https://doi.org/10.3928/01477447-20180213-05>.
- Foster, D., J. Williams, A. J. Forte, et al. n.d. "Application of Ice for Postoperative Total Knee Incisions – Does This Make Sense? A Pilot Evaluation of Blood Flow Using Fluorescence Angiography." *Cureus* 11 (7): e5126. <https://doi.org/10.7759/cureus.5126>.
- Freccero, D. M., P. Van Steyn, P. M. N. Joslin, et al. 2022. "Continuous Femoral Nerve Block Reduces the Need for Manipulation Following Total Knee Arthroplasty." *JBJS Open Access* 7 (3): e21.00155. <https://doi.org/10.2106/JBJS.OA.21.00155>.
- Gasparini, G., F. Familiari, and F. Ranuccio. 2013. "Patellar Malalignment Treatment in Total Knee Arthroplasty." *Joints* 1 (1): 10–17.
- Ghritlahre, A. K., G. S. Sandhu, S. Gupta, R. K. Garg, H. Dhindsa, and A. Gill. 2024. "Effect of Early versus Delayed Enteral Feeding in Case of Perforation Peritonitis." *Int J Life Sci Biotechnol Pharma Res* 13:582–88. [https://doi.org/10.69605/ijlbr\\_13.8.2024.100](https://doi.org/10.69605/ijlbr_13.8.2024.100).
- Harato, K., H. Tanikawa, Y. Morishige, K. Kaneda, and Y. Niki. 2016. "What Are the Important Surgical Factors Affecting the Wound Healing after Primary Total Knee Arthroplasty?" *J Orthop Surg* 11:7. <https://doi.org/10.1186/s13018-016-0340-y>.
- Helito, C. P., D. K. Bueno, P. N. Giglio, M. B. Bonadio, J. R. Pécora, and M. K. Demange. 2017. "NEGATIVE-PRESSURE WOUND THERAPY IN THE TREATMENT OF COMPLEX INJURIES AFTER TOTAL KNEE ARTHROPLASTY." *Acta Ortop Bras* 25 (2): 85–88. <https://doi.org/10.1590/1413-785220172502169053>.
- Ipach, I., F. Mittag, J. Lahrmann, B. Kunze, and T. Kluba. 2011. "Arthrofibrosis after TKA - Influence Factors on the Absolute Flexion and Gain in Flexion after Manipulation under Anaesthesia." *BMC Musculoskelet Disord* 12 (1): 184. <https://doi.org/10.1186/1471-2474-12-184>.
- Jaiswal, P. K., J. R. Perera, W. Khan, and S. G. Rao. 2012. "Treating Stiffness After Total Knee Arthroplasty: A Technical Note and Preliminary Results." *Open Orthop J* 6:276–80. <https://doi.org/10.2174/1874325001206010276>.
- Jiang, Y., A. A. Trotsyuk, S. Niu, et al. 2023. "Wireless, Closed-Loop, Smart Bandage with Integrated Sensors and Stimulators for Advanced Wound Care and Accelerated Healing." *Nat Biotechnol* 41 (5): 652–62. <https://doi.org/10.1038/s41587-022-01528-3>.
- Jurecka, A., M. Papież, P. Skucińska, and A. Gądek. 2021. "Evaluating the Effectiveness of Soft Tissue Therapy in the Treatment of Disorders and Postoperative Conditions of the Knee Joint-A Systematic Review." *J Clin Med* 10 (24): 5944. <https://doi.org/10.3390/jcm10245944>.
- Khalefa, M. A., L. K. Smith, and R. Ahmad. 2020. "The Use of Tissue Adhesive as an Adjunct to Wound Closure in Knee Arthroplasty Does Not Reduce Wound Ooze." *Knee Surg Relat Res* 32 (1): 57. <https://doi.org/10.1186/s43019-020-00073-0>.
- Khlopas, A., N. Sodhi, H. K. Anis, J. Ehiorobo, A. A. Sultan, and M. A. Mont. 2019. "A Systematic Review of Suture Technologies in Total Knee Arthroplasty." *Surg Technol Int* 34:391–96.

- Kong, X., M. Yang, Z. Cao, J. Chen, W. Chai, and Y. Wang. 2020. "Tissue Adhesive for Wound Closure in Enhanced-Recovery Total Hip Arthroplasty: A Prospective, Randomized and Controlled Study." *BMC Musculoskelet Disord* 21:178. <https://doi.org/10.1186/s12891-020-03205-5>.
- Krebs, V. E., R. K. Elmallah, A. Khlopas, et al. 2018. "Wound Closure Techniques for Total Knee Arthroplasty: An Evidence-Based Review of the Literature." *J Arthroplasty* 33 (2): 633–38. <https://doi.org/10.1016/j.arth.2017.09.032>.
- Kulkarni, S., M. Goodbun, M. Chowdhury, and P. W. Stather. 2025. "Dermabond Prineo: A Systematic Review and Meta-Analysis." *J Wound Care* 34 (3): 220–26. <https://doi.org/10.12968/jowc.2023.0024>.
- Kurdi, M., A. Moukhtar, M. Elkholy, H. Alwassia, M. Bamehriz, and M. G. Khirallah. 2023. "Delayed vs. Early Enteral Feeding after Repair of Congenital Recto-Vestibular Fistula: The Effect on Perineal Wound Healing." *Front Pediatr* 10:994249. <https://doi.org/10.3389/fped.2022.994249>.
- Lauerman, M., O. Kolesnik, H. Park, et al. 2018. "Definitive Wound Closure Techniques in Fournier's Gangrene." *Am Surg* 84 (1): 86–92. <https://doi.org/10.1177/000313481808400127>.
- Lee, G. W., W. K. Kwak, and K. B. Lee. 2021. "Comparison of 2-Octyl Cyanoacrylate Skin Adhesive and Interrupted Polypropylene Sutures for Wound Closure in Total Ankle Arthroplasty." *J Orthop Surg* 16 (1): 636. <https://doi.org/10.1186/s13018-021-02791-x>.
- Lee, H. G. 2024. "Is Liquid Skin Adhesive Safe and Feasible for Skin Closure in Single-Port Laparoscopic Appendectomy?" *J Minim Invasive Surg* 27 (2): 74–75. <https://doi.org/10.7602/jmis.2024.27.2.74>.
- Li, E., W. Niu, T. Lu, et al. 2020. "A Comparison between Barbed Suture and Conventional Suture in Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *Arthroplasty* 2 (1): 8. <https://doi.org/10.1186/s42836-020-00028-6>.
- Li, L., Q. Shao, W. He, T. Wang, and F. Wang. 2023. "Close Orthopedic Surgery Skin Incision with Combination of Barbed Sutures and Running Subcuticular Suturing Technique for Less Dermal Tension Concentration: A Finite Element Analysis." *J Orthop Surg* 18 (1): 333. <https://doi.org/10.1186/s13018-023-03755-z>.
- Liu, P., X. Mu, Q. Zhang, Z. Liu, W. Wang, and W. Guo. 2020. "Should Compression Bandage Be Performed after Total Knee Arthroplasty? A Meta-Analysis of Randomized Controlled Trials." *J Orthop Surg* 15:52. <https://doi.org/10.1186/s13018-019-1527-9>.
- Liu, T., Y. Tao, R. Zhao, et al. 2024. "Comparison of the Safety and Efficacy of Three Superficial Skin Closure Methods for Multi-Layer Wound Closure in Total Knee Arthroplasty: A Multicenter, Prospective, Randomized Controlled Trial." *Arthroplasty* 6:51. <https://doi.org/10.1186/s42836-024-00271-1>.
- Lu, X., L. Zhong, X. Cao, J. Liu, J. Chen, and D. Guo. 2021. "Wound Closure Position in Total Knee Arthroplasty: Flexion versus Extension—a Meta-Analysis of Randomized Controlled Trials." *Arch Orthop Trauma Surg* 141 (11): 1971–82. <https://doi.org/10.1007/s00402-021-03835-x>.
- Luo, X., W. Zhang, P. Yan, et al. 2020. "Skin Closure Tape and Surgical Staples in Primary Total Knee Arthroplasty: A Systematic Review and Meta-Analysis." *BioMed Res Int* 2020 (1): 4827617. <https://doi.org/10.1155/2020/4827617>.
- Maniar, R. N., B. Mody, H. M. Wakankar, et al. 2023. "Expert Consensus on Best Practices for Optimal Wound Closure in Total Knee Arthroplasty: A STRIDE Initiative for Orthopedic Surgeons of India." *Indian J Orthop* 57 (12): 2066–72. <https://doi.org/10.1007/s43465-023-00922-6>.
- Mau-Moeller, A., M. Behrens, S. Finze, S. Bruhn, R. Bader, and W. Mittelmeier. 2014. "The Effect of Continuous Passive Motion and Sling Exercise Training on Clinical and Functional Outcomes Following Total Knee Arthroplasty: A Randomized Active-Controlled Clinical Study." *Health Qual Life Outcomes* 12:68. <https://doi.org/10.1186/1477-7525-12-68>.
- Medeiros, I., S. Pascoal, F. Castro, J. P. Macedo, O. Lopes, and J. Pereira. 2024. "Tissue Adhesives for Closure of Intraoral Surgical Incisions: A Systematic Review and Meta-Analysis." *Int Wound J* 21 (10): e70047. <https://doi.org/10.1111/iwj.70047>.
- Nambi Gowri, K., and M. W. King. 2023. "A Review of Barbed Sutures—Evolution, Applications and Clinical Significance." *Bioengineering* 10 (4): 419. <https://doi.org/10.3390/bioengineering10040419>.
- Naylor, B. H., J. M. Tarazi, H. S. Salem, S. F. Harwin, and M. A. Mont. 2023. "Wound Management Following Total Knee Arthroplasty: An Updated Review." *J Knee Surg* 36 (3): 274–83. <https://doi.org/10.1055/s-0041-1731740>.
- Nepal, S., P. Ruangsombon, P. Udomkiat, and A. Unnanuntana. 2020. "Cosmetic Outcomes and Patient Satisfaction Compared between Staples and Subcuticular Suture Technique for Wound Closure after Primary Total Knee Arthroplasty: A Randomized Controlled Trial." *Arch Orthop Trauma Surg* 140 (9): 1255–63. <https://doi.org/10.1007/s00402-020-03479-3>.

- Newman, J. T., S. J. Morgan, G. V. Resende, A. E. Williams, E. M. Hammerberg, and M. R. Dayton. 2011. "Modality of Wound Closure after Total Knee Replacement: Are Staples as Safe as Sutures? A Retrospective Study of 181 Patients." *Patient Saf Surg* 5:26. <https://doi.org/10.1186/1754-9493-5-26>.
- Nguyen, H. M., T. T. Ngoc Le, A. T. Nguyen, H. N. Thien Le, and T. T. Pham. n.d. "Biomedical Materials for Wound Dressing: Recent Advances and Applications." *RSC Adv* 13 (8): 5509–28. <https://doi.org/10.1039/d2ra07673j>.
- Nifontova, G., S. Safaryan, Y. Khristidis, et al. 2024. "Advancing Wound Healing by Hydrogel-Based Dressings Loaded with Cell-Conditioned Medium: A Systematic Review." *Stem Cell Res Ther* 15:371. <https://doi.org/10.1186/s13287-024-03976-x>.
- Park, Y. H., J. S. Choi, J. W. Choi, and H. J. Kim. 2021. "Incidence and Risk Factor of Allergic Contact Dermatitis to 2-Octyl Cyanoacrylate and n-Butyl Cyanoacrylate Topical Skin Adhesives." *Sci Rep* 11 (1): 23762. <https://doi.org/10.1038/s41598-021-03319-3>.
- Peng, X., X. Zhang, T. Cheng, M. Cheng, and J. Wang. 2015. "Comparison of the Quadriceps-Sparing and Subvastus Approaches versus the Standard Parapatellar Approach in Total Knee Arthroplasty: A Meta-Analysis of Randomized Controlled Trials." *BMC Musculoskelet Disord* 16:327. <https://doi.org/10.1186/s12891-015-0783-z>.
- Rahbari, H., M. Ahmadi, M. A. Doreh, S. Mahmoudi, P. Ghaemmaghami, and A. Fereidouni. 2025. "Comparison of Surgical Wound Infection and Dehiscence Following the Use of Two Methods of Nylon Sutures and Skin Staples in Staples in Diabetic Mellitus Patients Undergoing Total Knee Arthroplasty Surgery: A Randomized Clinical Trial Study." *BMC Musculoskelet Disord* 26:70. <https://doi.org/10.1186/s12891-024-08263-7>.
- Romanini, E., G. A. Zanolli, T. Ascione, et al. 2024. "Barbed Sutures and Skin Adhesives Improve Wound Closure in Hip and Knee Arthroplasty." *Knee Surg Sports Traumatol Arthrosc Off J ESSKA* 32 (2): 303–10. <https://doi.org/10.1002/ksa.12055>.
- Roumeliotis, L., and N. M. Graham. 2019. "Barbed Suture and Glue in Skin Closure during Lower Limb Arthroplasty: Reduced Delayed Discharge Due to Wound Exudate." *J Wound Care* 28 (11): 784–89. <https://doi.org/10.12968/jowc.2019.28.11.784>.
- Rousseau, T., C. Plomion, and K. Sandy-Hodgetts. 2022. "An Advanced Transparent Hydropolymer Wound Dressing for Undisturbed Post-Op Management of Surgical Wounds Following Hip and Knee Replacement: A Prospective Observational Series." *Int Wound J* 19 (6): 1456–62. <https://doi.org/10.1111/iwj.13742>.
- Rr, Y., M. Aj, S. Ls, F. Aw, M. Ma, and M. Al. 2019. "Impact of Early Mobilization on Length of Stay after Primary Total Knee Arthroplasty." *Ann Transl Med* 7 (4). <https://doi.org/10.21037/atm.2019.02.02>.
- Sanz-Ruiz, P., J. R. Caeiro-Rey, J. C. Martínez-Pastor, J. L. Martín-Alguacil, A. Murcia-Asensio, and J. Moreta. 2025. "Consensus Document on the Management of Wound Closure in Orthopaedic Surgery." *EFORT Open Rev* 10 (2): 82–94. <https://doi.org/10.1530/EOR-24-0002>.
- Schwarz, E. M., J. Parvizi, T. Gehrke, et al. 2019. "2018 International Consensus Meeting on Musculoskeletal Infection: Research Priorities from the General Assembly Questions." *J Orthop Res Off Publ Orthop Res Soc* 37 (5): 997–1006. <https://doi.org/10.1002/jor.24293>.
- Shah, D., and A. Shah. 2024. "TRIVECTOR ARTHROTOMY FOR TOTAL KNEE ARTHROPLASTY: A 'GATEWAY' FOR ALL KNEES." *Orthop Proc* 106-B (SUPP\_20): 16–16. <https://doi.org/10.1302/1358-992X.2024.20.016>.
- Silva, J. R., B. Burger, C. M. C. Kühl, T. Candreva, M. B. P. dos Anjos, and H. G. Rodrigues. 2018. "Wound Healing and Omega-6 Fatty Acids: From Inflammation to Repair." *Mediators Inflamm* 2018:2503950. <https://doi.org/10.1155/2018/2503950>.
- Silva, P. H. S. da, C. E. B. Lopes, L. B. Stallmach, et al. 2023. "Comparison of Different Pneumorrhaphy Methods after Partial Pulmonary Lobectomy in Dogs." *Anim Open Access J MDPI* 13 (17): 2732. <https://doi.org/10.3390/ani13172732>.
- Siqueira, M. B., D. Ramanathan, A. K. Klika, C. A. Higuera, and W. K. Barsoum. 2016. "Role of Negative Pressure Wound Therapy in Total Hip and Knee Arthroplasty." *World J Orthop* 7 (1): 30–37. <https://doi.org/10.5312/wjo.v7.i1.30>.
- Snyder, M. A., B. P. Chen, A. Hogan, and G. W. J. Wright. 2021. "Multilayer Watertight Closure to Address Adverse Events From Primary Total Knee and Hip Arthroplasty: A Systematic Review of Wound Closure Methods by Tissue Layer." *Arthroplasty Today* 10:180-189.e7. <https://doi.org/10.1016/j.artd.2021.05.015>.
- Song, Q. C., D. Li, Y. Zhao, et al. 2022. "Negative Pressure Wound Therapy Reduces the Incidence of Postoperative Wound Dehiscence and Surgical Site Infections after Total Knee Arthroplasty in Patients with Obesity." *Medicine (Baltimore)* 101 (27): e29641. <https://doi.org/10.1097/MD.00000000000029641>.

- Su, S., J. He, C. Wang, F. Gao, D. Zhong, and P. Lei. 2022. "A New Dressing System Reduces the Number of Dressing Changes in the Primary Total Knee Arthroplasty: A Randomized Controlled Trial." *Front Surg* 9. <https://doi.org/10.3389/fsurg.2022.800850>.
- Sundaram, K., N. S. Piuizzi, B. M. Patterson, K. L. Stearns, V. E. Krebs, and M. A. Mont. 2020. "Skin Closure with 2-Octyl Cyanoacrylate and Polyester Mesh after Primary Total Knee Arthroplasty Offers Superior Cosmetic Outcomes and Patient Satisfaction Compared to Staples: A Prospective Trial." *Eur J Orthop Surg Traumatol Orthop Traumatol* 30 (3): 447–53. <https://doi.org/10.1007/s00590-019-02591-4>.
- Sundresh, N. J., M. Devagi, and D. Gopikrishna. 2018. "Comparison between Sutures and Staplers-Which Is Better for Laparotomy Wound Closure." *J Med Sci Clin Res* 6:930–35.
- Swain, D., and A. Gupta. 2015. "Interfacial Growth during Closure of a Cutaneous Wound: Stress Generation and Wrinkle Formation." *Soft Matter* 11 (32): 6499–6508. <https://doi.org/10.1039/c5sm01135c>.
- Tarazi, J. M., B. H. Naylor, H. S. Salem, and M. A. Mont. 2020. "The Use of New Technology for Wound Management Following Total Knee Arthroplasty: Implications for the COVID-19 Crisis." *Surg Technol Int* 37:291–98.
- Templeton, M. M., A. I. Krebs, K. H. Kraus, and C. S. Hedlund. 2015. "Ex Vivo Biomechanical Comparison of V-Loc 180® Absorbable Wound Closure Device and Standard Polyglyconate Suture for Diaphragmatic Herniorrhaphy in a Canine Model." *Vet Surg* 44 (1): 65–69. <https://doi.org/10.1111/j.1532-950X.2014.12201.x>.
- Teo, B. J. X., W. Yeo, H. C. Chong, and A. H. C. Tan. 2018. "Surgical Site Infection after Primary Total Knee Arthroplasty Is Associated with a Longer Duration of Surgery." *J Orthop Surg* 26 (2): 2309499018785647. <https://doi.org/10.1177/2309499018785647>.
- Thompson, R., D. Novikov, Z. Cizmic, et al. 2019. "Arthrofibrosis after Total Knee Arthroplasty: Pathophysiology, Diagnosis, and Management." *Orthop Clin* 50 (3): 269–79.
- Wang, S., J. Xia, Y. Wei, J. Wu, and G. Huang. 2014. "Effect of the Knee Position during Wound Closure after Total Knee Arthroplasty on Early Knee Function Recovery." *J Orthop Surg* 9:79. <https://doi.org/10.1186/s13018-014-0079-2>.
- Yuan, F. Z., J. Y. Zhang, D. Jiang, and J. K. Yu. 2019. "Quadriceps-Sparing versus Traditional Medial Parapatellar Approaches for Total Knee Arthroplasty: A Meta-Analysis." *BMC Musculoskelet Disord* 20:117. <https://doi.org/10.1186/s12891-019-2482-7>.
- Yuenyongviwat, V., K. Iamthanaporn, T. Hongnarak, and B. Tangtrakulwanich. 2016. "A Randomised Controlled Trial Comparing Skin Closure in Total Knee Arthroplasty in the Same Knee: Nylon Sutures versus Skin Staples." *Bone Jt Res* 5 (5): 185–90. <https://doi.org/10.1302/2046-3758.55.2000629>.
- Yusuf, K. 2022. "The Role of TNF-Alpha in the Wound Healing Process: Molecular and Clinical Perspectives - A Systematic Literature Review." *J RSMH Ptb* 3 (2): 222–28. <https://doi.org/10.37275/jrp.v3i2.53>.
- Zarepour, A., B. Gok, Y. Budama-Kilinc, A. Khosravi, S. Iravani, and A. Zarrabi. 2024. "Bacterial Nanocelluloses as Sustainable Biomaterials for Advanced Wound Healing and Dressings." *J Mater Chem B* 12 (48): 12489–507. <https://doi.org/10.1039/d4tb01024h>.
- Zhang, L., Y. Ning, C. Yang, and T. He. 2025. "Limiting Tourniquet Use during Total Knee Arthroplasty Improves Short-Term Postoperative Outcomes in Patients with Hypertension." *Front Surg* 12:1535662. <https://doi.org/10.3389/fsurg.2025.1535662>.
- Zhang, W., J. Xie, and A. Zeng. 2022. "The Origin and Development of Interrupted Subcuticular Suture: An Important Technique for Achieving Optimum Wound Closure." *Dermatol Surg* 48 (6): 619–24. <https://doi.org/10.1097/DSS.0000000000003437>.
- Zhang, X., Y. Huang, T. Luo, et al. 2024. "Advanced Wound Healing and Scar Reduction Using an Innovative Anti-ROS Polysaccharide Hydrogel with Recombinant Human Collagen Type III." *ACS Appl Mater Interfaces* 16 (38): 50305–20. <https://doi.org/10.1021/acsami.4c09890>.
- Zhu, Y., Y. Li, C. Yan, X. Du, Z. Xing, and P. Chen. 2017. "[Influence of Lateral Retinacular Release on Anterior Knee Pain Following Total Knee Arthroplasty]." *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi Zhongguo Xiu fu Chongjian Waikē Zazhi Chin J Reparatve Reconstr Surg* 31 (5): 541–46. <https://doi.org/10.7507/1002-1892.201701052>.